

**Hinweis/Note:****PENTAPHARM**

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.  
The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

# Ecarin

**Description:** Ecarin is a metalloprotease isolated from the venom of the saw-scaled viper (*Echis carinatus*) that activates prothrombin. Ecarin does not affect other clotting factors. The action of ecarin on prothrombin is independent of calcium, phospholipids and factor V. Moreover, Ecarin, in contrast to factor Xa, activates the acarboxy form of prothrombin present in plasma of patients undergoing oral anticoagulant therapy with vitamin K antagonists.

**Application:** Determination of prothrombin levels in patients undergoing anticoagulant therapy. Based on the interaction of hirudin with meizothrombin a simple ecarin clotting test for the monitoring of hirudin levels has been developed (Nowak and Bucha, 1993). Due to its action on acarboxy prothrombin, ecarin can be used for the determination of acarboxy prothrombin in the supernatant of barium sulfate treated patient plasma where normal carboxylated prothrombin has been removed by adsorption. Ecarin can be used for the detection of low prothrombin concentrations in plasma fractions for quality- and in-process control purposes. Ecarin, used in conjunction with the phospholipid- and calcium ion-dependent prothrombin activator Textarin® from *Pseudonaja textilis* venom, forms a highly sensitive and specific test system for lupus anticoagulant (Triplett et al., 1993).

**Origin:** *Echis carinatus* snake venom **MW:** 55'000 to 60'000

**Reconstitution:** Reconstitute vial with HEPES-buffered saline (0.2 M) containing 0.025 M calcium chloride. Recommended final ecarin concentration: 4 - 8 U/ml.

**Storage:** May be used by the expiry date given on the label when stored unopened, protected from moisture, in the dark, 2° - 8°C. Avoid contamination of the reagents by micro-organisms. Shipment of product does not require cooling during the time of transportation.

**Stability after reconstitution:**

+2°-8°C	7 days
- 20°C	6 months

**References:** Triplett DA, Stocker KF, Unger GA, Barna LK. The Textarin/Ecarin ratio: a confirmatory test for lupus anticoagulants. *Thromb Haemost* 1993; 70: 925-31.

Nowak G. Monitoring of the action of antithrombin agents by ecarin clotting time. In: *New anticoagulants for the cardiovascular patient*. Pifarré R, ed. Philadelphia: Hanley & Belfus 1997; 539-48.

Pöttsch B, Madlener K, Seelig C, Riess CF, Greinacher A, Müller-Berghaus G. Monitoring of r-hirudin anticoagulation during cardiopulmonary bypass assessment of the whole blood ecarin clotting time. *Thromb Haemost* 1997; 77: 920-25.

Nowak G. Clinical Monitoring of Hirudin and Direct Thrombin Inhibitors. *Sem. Thromb Haemost* 2001; 27: 537-41.

**Package size:** vials contains 50 U

**Code:** 116-01

FOR RESEARCH USE ONLY. NOT FOR HUMAN USE OR DRUG USE.

Registered as  
DSM Nutritional Products Ltd Branch Pentapharm  
Dornacherstrasse 112, CH-4147 Aesch BL/Switzerland  
Phone: +41 61 706 48 48, Fax: +41 61 706 48 00, [www.pentapharm.com](http://www.pentapharm.com)

**Distributed by:**

**LOXO** GMBH  
IMMUNOLOGIE • MOLEKULARBIOLOGIE  
BIOCHEMIE • PRODUKTE UND SYSTEME

69215 Dossenheim, Postfach 1130  
Tel.: +49 6221 868023  
Fax: +49 6221 8680255  
E-Mail: [info@loxo.de](mailto:info@loxo.de)  
Internet: [www.loxo.de](http://www.loxo.de)