

Monoclonal antibody against human CD31 PE conjugated

Product Nos. ADG5018 and ADG5018L

Description

CD31 (platelet endothelial cell adhesion molecule-1, PECAM-1) is an inhibitory coreceptor involved in regulation of T cell and B cell signaling by a dual immunoreceptor tyrosine-based inhibitory motif (ITIM) that upon associated kinases-mediated phosphorylation provide docking sites for protein-tyrosine phosphatases. CD31 is expressed ubiquitously within the vascular compartment and is located mainly at junctions between adjacent cells. N-terminal Ig-like domain of CD31 is responsible for its homophilic binding, which plays an important role in cell-cell interactions. CD31 is a multifunctional molecule with diverse roles in modulation of integrin-mediated cell adhesion, transendothelial migration, angiogenesis, apoptosis, negative regulation of immunoreceptor signaling, autoimmunity, macrophage phagocytosis, IgE-mediated anaphylaxis and thrombosis. It is one of key regulatory molecules in vascular system.

Properties

The monoclonal antibody ADG5018/L (clone MEM-05) is a murine monoclonal antibody, subclass IgG₁. The antibody has been purified from ascites by protein-A affinity chromatography, Purity > 95% (by SDS-PAGE).

The antibody reacts with CD31 (PECAM-1), a 130-140 kDa type I transmembrane glycoprotein expressed on monocytes, platelets, granulocytes, endothelial cells and stem cells of the myeloid lineage.

Conjugation

The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Presentation

Vial containing 500 µl (ADG5018) or 2 ml (ADG5018L) of purified antibody in PBS containing 1% BSA and 0.09% sodium azide (pH 7.2). The IgG concentration is 1 mg/ml. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at 4°C. Avoid prolonged exposure to light. The reagent is stable until the expiry date stated on the vial label.

Applications

Flow Cytometry

References

Prager E, Staffler G, Majdic O, Saemann M, Godar S, Zlabinger G, Stockinger H.: Induction of hyporesponsiveness and impaired T lymphocyte activation by the CD31 receptor:ligand pathway in T cells. J Immunol. 2001 Feb 15;166(4):2364-71.

Distributed by:

LOXO GMBH

IMMUNOLOGIE • MOLEKULARBIOLOGIE
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
69221 Dossenheim

Tel +49 6221 868023
Fax +49 6221 8680255

www.loxo.de - info@loxoxo.de

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmBioMed will not be held responsible for patent infringement or other violations that may occur with the use of our products.