

## Monoclonal antibody against human CD42b APC conjugated

Product Nos. ADG5039 and ADG5039L

### Description

**CD42b** (GPIb $\alpha$ ) composes together with GPIb $\beta$ , GPIX and GPV the GPIb-IX-V receptor complex critical in the process of platelet-rich thrombus formation by tethering the platelet to a thrombogenic surface. CD42b binds to von Willebrand factor (VWF) exposed at a site of vascular injury, as well as to thrombin, coagulation factors XI and XII, high molecular weight kininogen, TSP-1, integrin Mac-1 and P-selectin. The extracellular domain of CD42b by its interactions also contributes to metastasis.

### Properties

The monoclonal antibody ADG5039/L (clone HIP1) is a murine monoclonal antibody, subclass IgG<sub>1</sub>. The antibody has been purified from by protein-G affinity chromatography, Purity > 95% (by SDS-PAGE).

The antibody recognizes a glycoprotein, heterodimer (GPIb $\alpha$ ) which contains an  $\alpha$  chain (CD42b, 145 kDa) and a  $\beta$  chain (CD42c, 22 kDa) linked by a disulfid bond forming a noncovalent complex with CD42a (GPIX) and d. This antigen is expressed on platelets and megakaryocytes. The GPIb/IX complex (CD42a-d) serves as a receptor for von Willebrand Factor (vWF) and thrombin, and mediates adhesion of platelets to subendothelium of damaged vascular walls. The absence of the CD42 complex leads to the Bernard-Soulier Syndrome (BSS). HIP1 McAv inhibits the ristocetin-dependent binding of vWF to platelets, ristocetin-induced platelets agglutination, and partially inhibits collagen-induced aggregation

### Conjugation

The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

### Presentation

Vial containing 500  $\mu$ l (ADG5039) or 2 ml (ADG5039L) of purified antibody in PBS containing 1% BSA and 0.09% sodium azide (pH 7.2). The IgG concentration is 1 mg/ml. Spin the vial briefly before opening.

### Storage and Stability

Store the antibody at 4°C. Avoid prolonged exposure to light. The reagent is stable until the expiry date stated on the vial label.

### Applications

Flow Cytometry

### References

Leucocyte Typing IV. Knapp W et al. (Eds.), Oxford University Press (1989).

### Distributed by:

**LOXO** GMBH

IMMUNOLOGIE • MOLEKULARBIOLOGIE  
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48  
69221 Dossenheim

Tel +49 6221 868023

Fax +49 6221 8680255

www.loxo.de - info@loxox.de

### Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmBioMed will not be held responsible for patent infringement or other violations that may occur with the use of our products.