

Monoclonal antibody against human CD43 FITC conjugated

Product Nos. ADG5042 and ADG5042L

Description

CD43 (leukosialin, sialophorin) is a transmembrane mucin-like protein with high negative charge, expressed on the surface of most hematopoietic cells. CD43 contributes to a repulsive barrier that interferes with cellular adhesion, however, in certain cases also promotes leukocyte aggregation. By interaction with actin-binding proteins ezrin and moesin CD43 plays a regulatory role in remodeling T-cell morphology and regulates cell-cell interactions during lymphocyte traffic. CD43 signaling both enhances LFA-1 adhesiveness and counteracts LFA-1 induction via other receptors. Expression of CD43 causes induction of functionally active tumour suppressor p53 protein, but in case of p53 and ARF deficiency CD43 promotes tumour proliferation and viability. It appears to be an important modulator of leukocyte functions.

Properties

The monoclonal antibody ADG5042/L (clone MEM-59) is a murine monoclonal antibody, subclass IgG₁. The antibody has been purified from cell culture supernatant using Protein A affinity chromatography, Purity > 95% (by SDS-PAGE).

The antibody recognizes neuraminidase-sensitive epitope on CD43 (Leukosialin), a 95135 kDa type I transmembrane glycoprotein (mucin-type) which is involved in lymphocyte activation. CD43 is expressed by platelets and at high levels on the surface of all leukocytes; it is negative on resting B lymphocytes and erythrocytes.

Conjugation

The purified antibody is conjugated with Fluorescein-isothiocyanate (FITC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Presentation

Vial containing 500 µl (ADG5042) or 2 ml (ADG5042L) of purified antibody in PBS containing 1% BSA and 0.09% sodium azide (pH 7.2). The IgG concentration is 1 mg/ml. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at 4°C. Avoid prolonged exposure to light. The reagent is stable until the expiry date stated on the vial label.

Applications

Flow Cytometry

References

- 1.) Deckert, M. et al., Eur. J. Immunol. 22, 2943, 1992.
- 2.) Alvarado, M. et al., Eur. J. Immunol. 25, 1051, 1995.
- 3.) Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- 4.) Cermak, L. et al., J. Biol. Chem. 227, 7955, 2002.

Distributed by:

LOXO GMBH

IMMUNOLOGIE • MOLEKULARBIOLOGIE
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
69221 Dossenheim

Tel +49 6221 868023

Fax +49 6221 8680255

www.loxo.de - info@loxo.de

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmBioMed will not be held responsible for patent infringement or other violations that may occur with the use of our products.