

Monoclonal antibody against human CD45 FITC conjugated

Product Nos. ADG5050 and ADG5050L

Description

CD45 (LCA, leukocyte common antigen) is a receptor-type protein tyrosine phosphatase ubiquitously expressed in all nucleated hematopoietic cells, comprising approximately 10% of all surface proteins in lymphocytes. CD45 glycoprotein is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases. CD45 protein exists as multiple isoforms as a result of alternative splicing; these isoforms differ in their extracellular domains, whereas they share identical transmembrane and cytoplasmic domains. These isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. Besides the role in immunoreceptor signaling, CD45 is important in promoting cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis.

Properties

The monoclonal antibody ADG5050/L (clone MEM-28) is a murine monoclonal antibody, subclass IgG₁. The antibody has been purified from cell culture supernatant using protein-A affinity chromatography, Purity > 95% (by SDS-PAGE).

The antibody reacts with all alternative forms of human CD45 antigen (Leukocyte Common Antigen), a 180-220 kDa single chain type I transmembrane protein expressed at high level on all cells of hematopoietic origin, except erythrocytes and platelets.

Distributed by:



IMMUNOLOGIE • MOLEKULARBIOLOGIE
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
69221 Dossenheim

Tel +49 6221 868023
Fax +49 6221 8680255

www.loxo.de - info@loxode

Conjugation

The purified antibody is conjugated with Fluorescein-isothiocyanate (FITC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Presentation

Vial containing 500 µl (ADG5050) or 2 ml (ADG5050L) of purified antibody in PBS containing 1% BSA and 0.09% sodium azide (pH 7.2). The IgG concentration is 1 mg/ml. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at 4°C. Avoid prolonged exposure to light. The reagent is stable until the expiry date stated on the vial label.

Applications

Flow cytometry.

References

Horejsi V. et al., Folia Biol. (Praha) 34, 23 (1988)
Bazil Vet al., Immunogenetics 29, 202 (1989)
Leucocyte Typing IV. Knapp W. et al. (Eds), Oxford University Press (1989).

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmBioMed will not be held responsible for patent infringement or other violations that may occur with the use of our products.