

Monoclonal antibody against human CD45 APC conjugated

Product Nos. ADG5059 and ADG5059L

Description

CD45 (LCA, leukocyte common antigen) is a receptor-type protein tyrosine phosphatase ubiquitously expressed in all nucleated hematopoietic cells, comprising approximately 10% of all surface proteins in lymphocytes. CD45 glycoprotein is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases. CD45 protein exists as multiple isoforms as a result of alternative splicing; these isoforms differ in their extracellular domains, whereas they share identical transmembrane and cytoplasmic domains. These isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. Besides the role in immunoreceptor signaling, CD45 is important in promoting cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis.

Properties

The monoclonal antibody ADG5059/L (clone HI30) is a murine monoclonal antibody, subclass IgG₁. The antibody has been purified from ascites using protein-A affinity chromatography, Purity > 95% (by SDS-PAGE).

The antibody reacts with all alternative forms of human CD45 antigen (Leukocyte Common Antigen), a 180-220 kDa single chain type I transmembrane protein expressed at high level on all cells of hematopoietic origin, except erythrocytes and platelets and non-hematopoietic tissues and cells. CD45 is critically required for T and B cell antigen receptor-mediated activation.

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

Conjugation

The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Presentation

Vial containing 500 µl (ADG5059) or 2 ml (ADG5059L) of purified antibody in PBS containing 1% BSA and 0.09% sodium azide (pH 7.2). The IgG concentration is 1 mg/ml. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at 4°C. Avoid prolonged exposure to light. The reagent is stable until the expiry date stated on the vial label.

Applications

Flow cytometry

References

Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York
Kishihara K, *et al.* 1993. *Cell* 74:143.
Esser M, *et al.* 2001. *J. Virol.* 75:6173. (WB)
Friedman T, *et al.* 1999. *J. Immunol.* 162:5256. (IHC)
Rees LE, *et al.* 2003. *Clin. Exp. Immunol.* 134:497. (IF)

Distributed by:



IMMUNOLOGIE • MOLEKULARBIOLOGIE
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
69221 Dossenheim

Tel +49 6221 868023

Fax +49 6221 8680255

www.loxo.de - info@loxode

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmBioMed will not be held responsible for patent infringement or other violations that may occur with the use of our products.