

**Monoclonal antibody against outer surface lipoprotein BBA03
Borrelia burgdorferi [A03-2.4]
Product No. ADG0117L**

Description

Lyme disease is the most common vector-borne disease in North America and Europe. The causative agent *Borrelia burgdorferi* is a bacterium that is maintained in an enzootic cycle between *Ixodes* ticks and a large range of mammals. Linear plasmid lp54 is one of the most highly conserved and differentially expressed elements of the segmented genome of *Borrelia burgdorferi*. Several outer surface proteins encoded on the lp54 plasmid have been shown to be critical at specific times during the infectious cycle. Among these are the OspA/B, DbpA/B, BBA03, BBA52, and BBA64.

Properties

The monoclonal antibody ADG0117L (clone **A03-2.4**) is a murine monoclonal antibody, subclass IgG₁ recognizing an outer surface 20 kDa lipoprotein (BBA03). Mice were immunized with cell lysates of *Borrelia burgdorferi*. The antibody has been purified from cell culture supernatant using Protein G affinity chromatography.

Presentation

Screw capped vial containing 1 mg of purified antibody in PBS pH 7.4. The IgG concentration is given on the vial label. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at 2°-8°C. For long-term storage the antibody should be aliquoted and stored at -20°C or colder. It is recommended to avoid freeze-thaw cycles.

Applications

A. ELISA

The antibody can be used as capture antibody in ELISAs. An antibody concentration of 1-10 µg/ml is recommended.

B. Westernblot

The antibody is suitable for Western blot analysis, detecting native and recombinant lpLA7 following SDS-PAGE under reducing conditions. A primary antibody concentration of 1-10 µg/mL is recommended.

C. Immunocytochemistry

The antibody can be used for immunocytochemistry on paraformaldehyde fixed.

References

1. Competitive advantage of *Borrelia burgdorferi* with outer surface protein BBA03 during tick-mediated infection of the mammalian host. Bestor et al. *Infect. Immun.* 2012; 80(10):3501-3511
2. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. Fraser et al. *Nature* 1997; 390:580-586
3. Comprehensive spatial analysis of the *Borrelia burgdorferi* lipoproteome reveals a compartmentalization bias toward bacterial surface. Dowdell et al. *J. Bacteriol.* 2017; 199(6):e00658-16
4. Identification of 11 pH-regulated genes in *Borrelia burgdorferi* localizing to linear plasmids. Carroll et al. *Infect. Immun.* 2000; 68(12):6677-6684

Distributed by:

LOXO GMBH

IMMUNOLOGIE • MOLEKULARBIOLOGIE
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
69221 Dossenheim

Tel +49 6221 868023
Fax +49 6221 8680255

www.loxo.de - info@loxode

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.