IMMBIOMED

ImmBioMed GmbH & Co. KG Bergstrasse 85, DE-64319 Pfungstadt Phone: +49 6157 9799 685, Fax: +49 6157 86443

Monoclonal antibody against complement regulator-acquiring protein (HcpA) *Borrelia recurrentis* [Br-4]

Product No. ADG0162L

Description

Borrelia (B.) recurrentis, the causal agent of louse-borne relapsing fever is transmitted to humans via infected body lice. Infection with *B. recurrentis* has been achieved only in humans and is accompanied by a systemic inflammatory disease, multiple relapses of fever and massive spirochetemia. A key virulence factor of *B. recurrentis* is their potential to undergo antigenic variation. In addition, *B. recurrentis* express a surface lipoprotein, termed HcpA, which by exploiting host proteins factor H, CFHR-1, and plasmin(ogen) confers resistance to both, complement attack and opsonization and simultaneously acquires an increased potential to invade host tissues.

Properties

The monoclonal antibody ADG0162L (**Br-4**) is a murine monoclonal antibody, subclass IgG_{2b} recognizing HcpA of *Borrelia recurrentis* strain A1. Mice were immunized with rec. HcpA. The antibody has been purified from cell culture supernatant using Protein G affinity chromatography.

Presentation

Vial containing 1 mg purified antibody in PBS pH 7.4. The concentration is given on the vial label. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at $2^{\circ}-8^{\circ}$ C. For long-term storage the antibody should be aliquoted and stored at -20° C or colder. It is recommended to avoid freeze-thaw cycles.

Applications

A. ELISA

The antibody can be used as capture antibody in ELISAs. An antibody concentration of 1-10 $\mu g/ml$ is recommended.

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

B. Westernblot

The antibody is suitable for Western blot analysis, detecting native and recombinant HcpA following SDS-PAGE under reducing conditions. A primary antibody concentration of 1-10 μ g/mL is recommended.

C. Immunocytochemistry

The antibody can be used for immunocytochemistry on paraformaldehyde fixed spirochetes.

References

- 1. Borrelia recurrentis employs a novel multifunctional surface protein with anti-complement, anti-opsonic and invasive potential to escape innate immunity. Grosskinsky et al. *PloS One*. 2009; 4(3):e4858
- 2. Human complement regulators C4b-binding protein and C1 esterase inhibitor interact with a novel outer surface protein of *Borrelia recurrentis*. Grosskinsky et al. *PLoS Negl. Trop. Dis.* 2011; 4(6):e698
- 3. Immune evasion strategies of relapsing fever spirochetes. Rötterding and Kraiczy. *Front. Immunol.* 2020; 11:1560



IMMUNOLOGIE • MOLEKULARBIOLOGIE BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48 69221 Dossenheim Tel +49 6221 868023 Fax +49 6221 8680255 www.loxo.de - info@loxo.de