

Heparin Cofactor II Deficient Plasma

Product No. ADG208

Lot XXXXXX

Description

Heparin Cofactor II (HCII) Deficient Plasma is manufactured from pooled normal human plasma depleted of HCII using antibodies directed to Factor HCII. Only the highest quality citrated plasma is used as starting material.

Specification

Activity assay	Lot XXXXXX
PT (Staaggo Neoplasine C1+)	XXX sec
APTT (HemosIL APTT-SP)	XXX sec
Fibrinogen (clottable)	XXX g/L
Heparin Cofactor II (antigen by ELISA)	< XXX U/ml

Application

For research assays of Heparin Cofactor II antigen and functionality as a negative control and for dilution of standards and test samples.

Warning

The plasma is prepared from screened normal citrated human plasma that have been tested and found negative for HIV and Hepatitis by FDA-approved methods.

As no known test method can provide complete assurance that products derived from human blood will not transmit HBsAg, HCV, HIV-1, HIV-2 or other blood-borne pathogens, this reagent should be handled as recommended for any potentially infectious human specimen.

Presentation and Storage

Set of 5 clear, screw-capped vials each containing 1 mL of lyophilized plasma. Plasma contains 50 mM HEPES and stabilizers. Store lyophilized plasma at 2°-8°C.

Reconstitution

One vial of lyophilized product is reconstituted by addition of 1 mL of distilled water, employing gentle agitation for 10 minutes. Reconstituted plasma should be kept on ice and used within 4 hours of dissolution.

Distributed by:

LOXO GMBH

IMMUNOLOGIE • MOLEKULARBIOLOGIE
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
69221 Dossenheim

Tel +49 6221 868023

Fax +49 6221 8680255

www.loxo.de - info@loxox.de

Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.
The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

For research use only!

ADG208©08072019_Lot