

Monoclonal antibody against Outer surface protein B (OspB)
***Borrelia burgdorferi* [LA-25.1]**
 Product No. ADG0084L

Description

Lyme disease is the most common vector-borne disease in North America and Europe. The causative agent *Borrelia burgdorferi* is a bacterium that is maintained in an enzootic cycle between *Ixodes* ticks and a large range of mammals. The OspB gene is localized on the same 49-kb linear plasmid as OspA, and both genes are transcribed from a common promoter. *B. burgdorferi* upregulates OspA and OspB during entry into ticks. OspA and OspB contribute to the colonization of bacterium within the vector gut and facilitate survival of *B. burgdorferi* in ticks.

Properties

The monoclonal antibody ADG0084L (clone LA-25.1) is a murine monoclonal antibody, subclass IgG_{2b} recognizing OspB. Mice were immunized with cell lysates of *Borrelia burgdorferi*. The antibody has been purified from cell culture supernatant using Protein G affinity chromatography.

Presentation

Vial containing 1 mg purified antibody in PBS pH 7.4. The concentration is given on the vial label. Spin the vial briefly before opening.

Storage and Stability

Store the antibody at 2°-8°C. For long-term storage the antibody should be aliquoted and stored at -20°C or colder. It is recommended to avoid freeze-thaw cycles.

Applications

A. ELISA


The antibody can be used as capture antibody in ELISAs. An antibody concentration of 1-10 µg/ml is recommended.

B. Westernblot

The antibody is suitable for Western blot analysis, detecting native and recombinant OspB following SDS-PAGE under reducing conditions. A primary antibody concentration of 1-10 µg/mL is recommended.

C. Immunocytochemistry

The antibody can be used for immunocytochemistry on paraformaldehyde fixed spirochetes and for cryo-electron tomography.



References

1. Characterization of *Borrelia burgdorferi* associated antigens by monoclonal antibodies. Kramer et al. *Immunobiol.* 1990; 181:357-366
2. Selection of an escape variant of *Borrelia burgdorferi* by use of bactericidal monoclonal antibodies to OspB. Coleman et al. *Infect. Immun.* 1992; 60(8):3098-3104
3. Nanoscopic localization of surface-exposed antigens of *Borrelia burgdorferi*. Lemgruber et al. *Microsc. Microanal.* 2015; 21(3):680-688
4. Complement-mediated *in vitro* bactericidal activity of monoclonal antibodies reactive with outer-surface-protein OspB of *Borrelia burgdorferi*. Cevenini et al. *FEMS Microbiol. Lett.* 1992; 69(2):147-152
5. Molecular analysis of linear plasmid-encoded major surface proteins, OspA and OspB, of the Lyme disease spirochete *Borrelia burgdorferi*. Bergström et al. 1989; 3(4):479-486
6. Selective association of outer surface lipoproteins with the lipid rafts of *Borrelia burgdorferi*. Toledo et al. 2014; 5(2):e00899-14

Distributed by:

LOXO GMBH

IMMUNOLOGIE • MOLEKULARBIOLOGIE
 BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48
 69221 Dossenheim

Tel +49 6221 868023

Fax +49 6221 8680255

www.loxo.de - info@loxo.de

For research use only!

Hinweis/Note:

ADG0084L©24082022

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.