

## Monoclonal antibody against the 41-kDa flagellin of *Borrelia burgdorferi* [LA-10] Product No. ADG0092L

### Description

Lyme disease is the most common vector-borne disease in North America and Europe. The causative agent *Borrelia burgdorferi* is a bacterium that is maintained in an enzootic cycle between *Ixodes* ticks and a large range of mammals. The 41 kDa flagellin of *Borrelia burgdorferi* is a major component of periplasmic flagellar filament core and a good candidate for serodiagnosis in early stage of Lyme disease.

### Properties

The monoclonal antibody ADG0092L (clone LA-10) is a murine monoclonal antibody, subclass IgG<sub>2a</sub> recognizing flagellin. Mice were immunized with cell lysates of *Borrelia burgdorferi*. The antibody has been purified from cell culture supernatant using Protein G affinity chromatography.

### Presentation

Vial containing 1 mg purified antibody in PBS pH 7.4. The concentration is given on the vial label. Spin the vial briefly before opening.

### Storage and Stability

Store the antibody at 2°-8°C. For long-term storage the antibody should be aliquoted and stored at -20°C or colder. It is recommended to avoid freeze-thaw cycles.

### Applications

#### A. ELISA


The antibody can be used as capture antibody in ELISAs. An antibody concentration of 1-10 µg/ml is recommended.

#### B. Westernblot

The antibody is suitable for Western blot analysis, detecting native and recombinant IpLA7 following SDS-PAGE under reducing conditions. A primary antibody concentration of 1-10 µg/mL is recommended.

### C. Immunocytochemistry

The antibody can be used for immunocytochemistry on paraformaldehyde fixed.



### References

1. Characterization of *Borrelia burgdorferi* associated antigens by monoclonal antibodies. Kramer et al. *Immunobiol.* 1990; 181:357-366
2. The *Borrelia burgdorferi* flagellum-associated 41-kilodalton antigen (flagellin): molecular cloning, expression, and amplification of the gene. Wallich et al. *Infect. Immun.* 1990; 58(6) 1711-1719
3. Cellular immune reactivity to recombinant OspA and flagellin from *Borrelia burgdorferi* in patients with Lyme borreliosis. Complexity of humoral and cellular immune responses. Krause et al. *J. Clin. Invest.* 1992; 90(3):1077-1084
4. Antigenic variation and strain heterogeneity in *Borrelia* spp. Wilske et al. *Res. Microbiol.* 1992; 143(6):583-596
5. Evaluation of genetic divergence among *Borrelia burgdorferi* isolates by use of OspA, fla, HSP60, and HSP70 gene probes. Wallich et al. *Infect. Immun.* 1992; 60(11):4856-4866

#### Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

**Distributed by:**

**LOXO**  
GMBH  
IMMUNOLOGIE • MOLEKULARBIOLOGIE  
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48  
69221 Dossenheim  
Tel +49 6221 868023  
Fax +49 6221 8680255  
www.loxo.de - info@lox.de

For research use only!

ADG0092L©ADG24082022