

## Protein S Deficient Plasma

Product No. ADG253

Lot XXXXXX

### Description

Protein S Deficient Plasma is manufactured from pooled normal human plasma depleted of Protein S using antibodies directed to Protein S immobilized on agarose beads.

### Specification

Activity assay	Lot XXXXXX
PT (Siemens Innovin)	XXX sec
APTT (Dade Actin FSL)	XXX sec
Fibrinogen (clottable)	XXX g/L
Factor II (activity)	XXX U/ml
Factor V (activity)	XXX U/ml
Factor VII (activity)	XXX U/ml
Factor VIII (activity)	XXX U/ml
Factor IX (activity)	XXX U/ml
Factor X (activity)	XXX U/ml
Factor XI (activity)	XXX U/ml
Factor XII (activity)	XXX U/ml
Protein S (antigen) by ELISA	< XXX U/ml
Protein C (antigen) by ELISA	XXX U/ml

### Application

For research assays of Protein S antigen and functionality, as a negative control and for dilution of standards and test samples.

### Distributed by:



IMMUNOLOGIE • MOLEKULARBIOLOGIE  
BIOCHEMIE • PRODUKTE UND SYSTEME

Gerhart-Hauptmann-Str. 48  
69221 Dossenheim

Tel +49 6221 868023

Fax +49 6221 8680255

www.loxo.de - info@loxo.de

### Warning

The plasma is prepared from normal citrated human plasma collected from donors screened for CJD and which was tested and found negative for HBsAg, syphilis (RPR or T Pallidum-G-EIA) and antibodies to HIV and HCV and non-reactive for HIV-1 RNA and HCV RNA by FDA approved methods.

As no known test method can provide complete assurance that products derived from human blood will not transmit HBsAg, HCV, HIV-1, HIV-2 or other blood-borne pathogens, this reagent should be handled as recommended for any potentially infectious human specimen.

### Presentation

Set of 5 clear, screw-capped vials each containing 1 mL of lyophilized plasma. The plasma contains 50mM HEPES and stabilizers.

### Reconstitution and Stability

One vial of lyophilized product is reconstituted by addition of 1 mL of distilled water, employing gentle agitation for 15 minutes, at room temperature. Stability after reconstitution is 4 hours at ambient (18° -25°C), or 30 days at -20 °C.

### Storage

Prior to reconstitution, dried plasma should be stored at 2° - 8°C.

#### Hinweis/Note:

Der Packungsbeileger dient nur als erste Information. Der relevante Packungsbeileger liegt der Ware bei.

The datasheet is for information purposes only. The current datasheet will be enclosed with product shipment.

**For research use only!**